American Journal of Emergency Medicine 93 (2025) 86–93

Contents lists available at ScienceDirect

American Journal of Emergency Medicine

journal homepage: www.elsevier.com/locate/ajem

Emergency medicine updates: Cardiopulmonary resuscitation

Brit Long, MD [a,*], Michael Gottlieb, MD [b]

[a] Department of Emergency Medicine, University of Virginia, Charlottesville, VA, USA
[b] Department of Emergency Medicine, Rush University Medical Center, Chicago, IL, USA

ARTICLE INFO

Article history:
Received 22 February 2025
Received in revised form 21 March 2025
Accepted 22 March 2025
Available online xxxx

Keywords:
Cardiac arrest
Return of spontaneous circulation
Resuscitation
Cardiopulmonary resuscitation
Compressions
Monitoring
ROSC
CPR
Compression fraction

ABSTRACT

Introduction: Cardiac arrest is the loss of functional cardiac activity; emergency clinicians are integral in the management of this condition.
Objective: This paper evaluates key evidence-based updates concerning cardiopulmonary resuscitation (CPR).
Discussion: Cardiac arrest includes shockable rhythms (i.e., pulseless ventricular tachycardia and ventricular fibrillation) and non-shockable rhythms (i.e., asystole and pulseless electrical activity). The goal of cardiac arrest management is to achieve survival with a good neurologic outcome, in part by restoring systemic perfusion and obtaining return of spontaneous circulation (ROSC), while seeking to diagnose and treat the underlying etiology of the arrest. CPR includes high-quality chest compressions to optimize coronary and cerebral perfusion pressure. Chest compressions should be centered over the mid-sternum, with the compressor's body weight over the middle of the chest. A compression depth of 5–6 cm is recommended at a rate of 100–120 compressions per minute, while allowing the chest to fully recoil between each compression. Clinicians should seek to minimize any interruptions in compressions. When performed by bystanders, compression-only CPR may be associated with improved survival to hospital discharge when compared to conventional CPR with ventilations. However, in trained personnel, there is likely no difference with compression-only versus conventional CPR. Mechanical approaches for CPR are not associated with improved patient outcomes, including ROSC or survival with good neurologic function, but mechanical compression devices may be beneficial in select circumstances (e.g., few rescuers available, prolonged arrest/transport). Monitoring of chest compressions is not associated with improved ROSC, survival, or neurologic outcomes, but it can improve guideline adherence. Types of monitoring include real-time feedback, a CPR coach, end tidal CO_2, arterial line monitoring, regional cerebral tissue oxygenation, and point-of-care ultrasound.
Conclusions: An understanding of CPR literature updates can improve the ED care of patients in cardiac arrest.
Published by Elsevier Inc.

1. Introduction

Cardiac arrest is the loss of functional cardiac activity and systemic circulation and includes shockable rhythms (i.e., pulseless ventricular tachycardia [pVT] and ventricular fibrillation [VF]) and non-shockable rhythms (i.e., asystole and pulseless electrical activity). It has an annual incidence ranging between 55 and 113 per 100,000 population, with 180,000–450,000 patients experiencing out-of-hospital cardiac arrest (OHCA) in the United States annually [1-6]. Most cases of cardiac arrest in those older than 35 years are associated with a cardiovascular etiology [7,8]. Unfortunately, mortality is significant, with data from 2022 suggesting survival to hospital discharge for patients experiencing OHCA treated by emergency medical services (EMS) is approximately 9 %, while survival to hospital discharge with good functional status is 7 % [3].

Cardiopulmonary resuscitation (CPR) is integral to improving outcomes in those with cardiac arrest and seeks to obtain survival with good neurologic outcome, in part by restoring systemic circulation [9-15]. CPR includes high-quality chest compressions to optimize coronary and cerebral perfusion pressure [1-3]. This review is part of a series discussing cardiac arrest management in the emergency department (ED) and will focus on CPR. Other reviews in this series will cover cardiac arrest medications, airway management, defibrillation strategies, ultrasound, and extracorporeal membrane oxygenation.

* Corresponding author at: University of Virginia, Department of Emergency Medicine, 200 Jeanette Lancaster Way, Charlottesville, VA 22903, USA.
E-mail address: Brit.long@yahoo.com (B. Long).

https://doi.org/10.1016/j.ajem.2025.03.057
0735-6757/Published by Elsevier Inc.

Descargado para Anonymous User (n/a) en Cayetano Heredia Peruvian University de ClinicalKey.es por Elsevier en abril 11, 2025. Para uso personal exclusivamente. No se permiten otros usos sin autorización. Copyright ©2025. Elsevier Inc. Todos los derechos reservados.

SHOUT
SHARE
IT

2. Discussion

2.1. What is the optimal technique and location for hand placement during chest compressions?

Cardiac arrest should be assumed in a person who is unresponsive and has absent or gasping breathing with no palpable pulse. The pulse should be checked at the same time the breathing is assessed, though this should take no longer than 10 s [11,16,17]. Following the recognition of cardiac arrest, high-quality chest compressions should be started as soon as possible, as these increase coronary and cerebral perfusion pressure and improve the likelihood of return of spontaneous circulation (ROSC) and survival with good neurologic outcome [9-15,18-24].

Integral measures of high-quality chest compressions include ensuring adequate rate and depth of compressions, while minimizing the frequency and duration of interruptions (Table 1) [1,2,9-15,22-24]. Current guidelines suggest a compression target of 100–120 beats per minute (bpm), with a chest compression fraction (total compression time divided by total resuscitation time) of at least 60 % and minimizing interruptions [11-15]. Of note, higher chest compression fractions (80 %) should be targeted if possible [14]. Compressions outside of this 100–120 bpm range reduce the likelihood of ROSC and are associated with worse outcomes [13,14,25-27]. A compression depth of 5–6 cm is recommended, with full chest recoil between compressions (Fig. 1) [12-15,25-30]. This results in negative intrathoracic pressure, increasing coronary perfusion pressure and cardiac preload [30]. Compressions may be improved with the compressor slightly removing their hands from the chest at the end of each compression.

Patients in cardiac arrest should immediately be placed supine, which is the most optimal position for high-quality compressions, with a hard surface behind the patient that serves as a counterforce to the compressions if this will not delay initiation of compression, as soft surfaces such as a mattress can significantly reduce compression depth [13,14,31,32]. This may include use of a backboard or the bed placed in CPR mode [13]. For in-hospital arrest, the International Liaison Committee on Resuscitation (ILCOR) guidelines recommend against moving the patient to the floor for compressions due to a delay in initiating compressions and creating a more hazardous environment [13]. In general, a backboard is preferred, as a meta-analysis of 6 studies found use of a backboard led to a 2 mm increase in chest compression depth, while CPR performed on a mattress or moving the patient to the floor did not affect compression depth [33]. However, the ILCOR guidelines state that if healthcare systems have not introduced backboards for use in cardiac arrest, there is insufficient evidence justifying their cost and training staff for their use [13].

The compressor should be standing beside the bed or kneeling next to the patient. Appropriate positioning may require lowering the height of the patient bed/stretcher or the compressor using a step stool. Appropriate hand positioning is also essential. The compressor's heel of their hand should be over the lower portion of the sternum, with the other heel of the hand over the first hand (Fig. 2) [1,2,13,14]. This ensures that compressions target the left ventricle, which is typically located inferior to a line connecting the nipples [34]. When performing compressions, the rescuer's chest should be directly above their hands with their arms in full extension [11-15]. This allows the rescuer to use their body weight to perform compressions, rather than relying on upper extremity strength, which can result in more rapid fatigue. Compressions may also be performed with the patient in the prone position, where the compressions are performed in the midline of the back at the T7-T9 level [35].

Importantly, rescuers often tire after one minute, which results in decreasing compression quality and recoil, but this must be balanced with the necessity of reducing interruptions in compressions, which leads to decreased coronary and cerebral perfusion pressures, ultimately reducing the likelihood of ROSC and survival [1,2,12-15,27,36-44]. Thus, the compressor should be switched every 2 min during rhythm and pulse check, which should not exceed 10 s. If compressions are significantly interrupted for longer than 10 s, it can take up to 1 min of high-quality compressions to restore sufficient perfusion pressures [1,2,11-14]. However, if the compressor is unable to perform high-quality compressions prior to the full 2 min due to fatigue, another person able to perform compressions should take over. To minimize interruptions, the defibrillator may be charged while compressions are ongoing until the shock can be delivered, and following defibrillation, compressions should be resumed immediately without repeat rhythm analysis [12-14,45-47]. The compressor may also wear gloves and continue compressions during defibrillation [48]. Procedures such as obtaining intravenous (IV) or interosseous (IO) access or endotracheal intubation should be performed while compressions are ongoing.

Table 1
Optimizing chest compressions.

- Center compressions over the mid-sternum, with the compressor's body weight over the middle of the chest.
- Compress to a depth of 5–6 cm (2–2.3 in.) with each compression.
- Allow the chest to fully recoil between compressions.
- Target a rate of 100–120 compressions per minute.
- Minimize the frequency and duration of interruptions in high-quality compressions, while maintaining a chest compression fraction over 60 % (over 80 % is optimal).
- Place a rigid surface under the patient or activate bed CPR mode to improve chest compression quality if possible; avoid moving the patient from the hospital bed to the ground.

2.2. Which CPR technique is optimal, compression-only or conventional CPR, and does the setting of the cardiac arrest make a difference in the recommended technique (e.g., lay provider/out-of-hospital versus trained provider as part of a response team)?

Conventional CPR incorporates a 30:2 compression-ventilation cycle if there is no definitive airway. If there is a definitive airway established, a breath should be delivered every 6 s with continuous chest compressions [1,2,11-14]. Compression-only CPR focuses on providing high-quality compressions while avoiding interruptions such as respirations in those without a definitive airway in order to improve coronary and cerebral perfusion pressures, as well as potentially improve the likelihood of successful defibrillation in shockable rhythms (Table 2) [12-15,27,49-53].

The rationale behind compression-only CPR is that conventional CPR may decrease blood flow and cardiac venous return and reduce coronary and cerebral perfusion during the compression pauses for breaths while increasing intrathoracic pressures [12-15,27,49-55]. Compression-only CPR may be most helpful in the initial 4–6 min of the cardiac arrest, when myocardial and cerebral tissues are most sensitive to reduced vascular flow and hypoperfusion leads to worse outcomes [11,14,56,57].

The Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care (CPRECC), which are a joint publication from ILCOR, the American Heart Association (AHA), and the European Resuscitation Council (ERC), state that high-quality compressions alone may be used if a sole rescuer is present or rescuers are reluctant to perform ventilations mouth-to-mouth [1,2,9-14,58,59]. Guidelines also state that rescuers should not interrupt compressions to check for pulses or ROSC and should continue high-quality compressions until an automated external defibrillator is prepared to defibrillate, if emergency medical services (EMS) providers assume management of the patient, or if the patient awakens [1,2,9-14].

Randomized data suggest a trend towards improved survival to hospital discharge with bystander performed compression-only CPR compared with conventional CPR [52,60,61], and a 2017 Cochrane review of 3 randomized control trials (RCTs) and 1 cluster RCT evaluating untrained bystander compression-only CPR found improved survival to hospital discharge when compared to conventional CPR [57]. However,

Descargado para Anonymous User (n/a) en Cayetano Heredia Peruvian University de ClinicalKey.es por Elsevier en abril 11, 2025. Para uso personal exclusivamente. No se permiten otros usos sin autorización. Copyright ©2025. Elsevier Inc. Todos los derechos reservados.



Fig. 1. Chest compression action on the heart.
(Obtained from https://commons.wikimedia.org/wiki/File:CPR-heart-compression.png.)

among EMS personnel, there was no difference. Other meta-analyses published after this Cochrane review have also found no difference in survival to hospital discharge or ROSC when evaluating compression-only and conventional CPR performed by trained personnel [62–65].

Based on the current data, compression-only CPR is likely associated with improved survival to hospital discharge when performed by bystanders when compared to conventional CPR with a 30:2 compression-ventilation ratio. However, in trained personnel, there is likely no difference with compression-only versus conventional CPR. A major consideration is the quality of compressions and chest compression fraction, which are both linked to improved outcomes. If trained personnel are alone, compression-only CPR is a viable option until assistance arrives.

2.3. When should mechanical approaches for CPR be utilized?

There are several mechanical approaches to CPR (Table 3). The most commonly utilized devices are mechanical compression devices, such as the AutoPulse (Fig. 3) (Zoll Medical Corporation, Chelmsford, MA) and Lund University Cardiopulmonary Assist System (LUCAS) (Fig. 4) (Physio-Control/Jolife AB, Lund, Sweden) currently approved by the U.S. Food and Drug Administration [66-70]. These devices differ in several aspects, though they all provide automatic delivery of compressions, minimizing interruptions in compressions, eliminating the issue of decreased perfusion from compressor fatigue, and allowing rescuers to perform other interventions.



Fig. 2. Hand location for appropriate chest compressions.
(Obtained from https://commons.wikimedia.org/wiki/File:Chest-compression-hand-placement.jpg.)

The AutoPulse has a load-distributing band that is placed circumferentially around the chest, with a back the shortens and lengthens, providing compressions [66]. The LUCAS device utilizes a piston which moves up and down over the same site as a rescuer's hands [67]. This piston is mounted on a circumferential frame with a power source, with a board underneath the patient's back. Data from multiple meta-analyses suggest no improvement in ROSC, survival to hospital discharge, and short-term or long-term survival with mechanical compression devices compared to manual compressions [70-74]. Of note, these devices require positioning around the patient and then connecting, adjusting, and initiating the device, which can result in pauses in compressions. Thus, training with these devices is necessary, as well as a standardized team approach for application of the device [75-77].

Active compression-decompression CPR (ACD CPR) incorporates a suction device that converts passive chest wall recoil into active expansion by pulling on the chest with an approximately 20 kg of force or distance of 3 cm, which can enhance venous return [78-80]. However, in isolation ACD CPR does not create adequate negative intrathoracic pressures [81]. A meta-analysis of 10 RCTs and quasi-RCTs (majority OHCA) found no benefit in patient-oriented outcomes including immediate survival, survival at hospital discharge, and survival with good neurologic function with ACD CPR compared to standard CPR [81]. Patients undergoing ACD CPR had greater rates of skin trauma and ecchymosis, though there was no difference in sternal fractures, hemothorax, or pneumothorax [81]. Based on this, the AHA states there is insufficient evidence for the use of ACD CPR [2,11-14].

Table 2
CPR technique considerations.

- Compression-only CPR has been evaluated and is likely associated with improved survival to hospital discharge when performed by bystanders compared to conventional CPR.
- In trained personnel, there is likely no difference in compression-only versus conventional CPR.
- The quality of chest compressions and chest compression fraction remain integral to optimizing neurologic outcomes in cardiac arrest.

Descargado para Anonymous User (n/a) en Cayetano Heredia Peruvian University de ClinicalKey.es por Elsevier en abril 11, 2025. Para uso personal exclusivamente. No se permiten otros usos sin autorización. Copyright ©2025. Elsevier Inc. Todos los derechos reservados.

Table 3
Mechanical devices for CPR considerations.

- A variety of mechanical devices for CPR are available (e.g., LUCAS, AutoPulse), but their use is not associated with improved patient outcomes.
- Mechanical compression devices may be assisted in cases with prolonged CPR or transport, limited availability of rescuers, and during coronary catheterization or extracorporeal membrane oxygenation cannulation.

Inspiratory impedance threshold devices (ITD) are designed to produce negative intrathoracic pressure during active or passive chest wall decompression, which may increase venous return to the heart during CPR and improve myocardial perfusion [82-85]. An ITD incorporates a plastic appliance with a silicone diaphragm between the airway (facemask, supraglottic device, endotracheal tube) and bag ventilator, which allows for positive-pressure ventilation and exhalation but does not allow air flow into the airway during relaxation of the chest wall. ITDs can produce negative intrathoracic pressures in patients undergoing bag-mask or invasive airway, and they are often used with ACD CPR to create effective negative intrathoracic force [82-85]. The combination of ITD and ACD CPR has demonstrated improved hemodynamics and vascular flow in preclinical trials [86,87]. The ROC PRIMED RCT included 8718 patients with OHCA and compared ITD with a sham device [84]. Authors found no significant difference in neurologically intact survival, ROSC on arrival, survival to admission, or survival to discharge. However, a 2015 secondary analysis of patients with "acceptable" CPR found that patients with an ITD had improved neurologically intact survival compared with the sham device (7.2 % vs. 4.1 %, P = 0.006) [88]. A 2010 study evaluating ACD CPR with an ITD compared to standard CPR in 1653 patients with OHCA found improved survival to hospital discharge in the ACD plus ITD group (8.9 % vs 5.8 %, P = 0.015), as well as increased survival with good neurologic outcome (8.6 % vs 5.8 %, P = 0.029) [89]. A meta-analysis found ITD was not associated with improved survival or neurologic outcomes, though when controlled for witnessed arrest and shorter response time, authors state ITD may improve ROSC [70]. Importantly, the AHA currently recommends against the routine use of ITD, though data suggest they may improve vascular flow [2,9-14].

Based on available evidence, mechanical approaches for CPR are not associated with improved patient outcomes, including survival, ROSC, or improved neurologic function. However, there are situations where mechanical compression devices such as the AutoPulse or LUCAS device may assist, including in cases of prolonged CPR or transport, limited availability of rescuers, and during coronary catheterization or extracorporeal membrane oxygenation cannulation [68-70].



Fig. 3. AutoPulse.
(Obtained from https://en.wikipedia.org/wiki/AutoPulse#/media/File:Mechanische_Reanimationshilfe.jpg.)

Fig. 4. LUCAS chest compression device.
(Obtained from https://en.wikipedia.org/wiki/LUCAS_device#/media/File:JASDF_Automatic_cardiopulmonary_resuscitation_device_at_Komaki_Air_Base_February_23,_2014.jpg.)

2.4. What is the evidence concerning head-up CPR?

Head-up CPR involves elevation of the head and chest during compressions (Table 4). This seeks to reduce intracranial pressure while maintaining mean arterial pressure to improve cerebral blood flow and cerebral perfusion pressure [90-92]. A 2019 study evaluating a prehospital bundle including head-up/chest-up CPR found this was safe and feasible during resuscitation [90]. A 2022 prospective registry study including 227 patients receiving head-up CPR combined with mechanical and/or ACD CPR with an ITD found a time-dependent association with ROSC. For every minute delay following the 911 phone call in applying the head-up CPR bundle, there was a 5.6 % reduction in likelihood of ROSC (between minutes 4–12) [91]. A second 2022 study evaluated head-up CPR combined with ACD CPR and ITD compared with conventional CPR. Authors found improved survival to discharge if this bundle was initiated within 18 min (odds ratio [OR] 1.88; 95 % confidence interval [CI] 1.03–3.44), with the greatest benefit within the first 9 min (OR 5.21; 95 CI% 1.74–15.59) [92]. However, there were several significant limitations, including no blinding or randomization, use of historical controls that may introduce confounding, and significant differences in the patient groups. In the 2022 study by Moore et al., the authors did not pre-establish time windows for their primary outcome, and they excluded four EMS systems, which increases the risk of selection bias [92]. The 2024 ILCOR Consensus Statement recommends that further trials are required evaluating the usefulness of head-up CPR due to the limited data [13].

Table 4
Head-up CPR considerations.

- Head-up CPR includes elevating the head and chest during compressions.
- Despite initial studies suggesting benefit, there are multiple limitations with the data (no blinding or randomization, use of historical controls that may introduce confounding, and significant differences in the patient groups).
- Further study is necessary prior to the routine use of head-up CPR.

Descargado para Anonymous User (n/a) en Cayetano Heredia Peruvian University de ClinicalKey.es por Elsevier en abril 11, 2025. Para uso personal exclusivamente. No se permiten otros usos sin autorización. Copyright ©2025. Elsevier Inc. Todos los derechos reservados.

2.5. What tools can be used to monitor chest compressions?

Chest compression quality is an integral component of cardiac arrest resuscitation, as inadequate CPR is associated with worse outcomes [11-14]. Even when performed by trained and experienced professionals, CPR quality varies and is often inadequate [11-14,93-95]. There are several means of monitoring chest compression rate and depth [11-14]. These include mechanical devices that provide real-time feedback of compression rate, depth, and chest recoil; use of a compression coach; end-tidal carbon dioxide ($EtCO_2$) measurements; diastolic blood pressure (DBP) measurement with invasive arterial pressure monitoring; regional cerebral tissue oxygenation (rSO_2); and point-of-care ultrasound (POCUS) (Table 5).

Mechanical devices can provide feedback to the compressor concerning compression rate and depth to ensure the performance of high-quality compressions. A 2020 systematic review conducted by ILCOR found no difference in patient outcomes with real-time feedback [15]. A 2023 systematic review found improved guideline compliance with chest compression depth and rate, but there was no difference in survival or ROSC [94]. A 2024 ILCOR systematic review found no improvement in patient outcomes [95]. However, authors did find that several aspects of CPR quality improved with real-time feedback provided by the device. The ILCOR 2024 Consensus statement recommends use of real-time audiovisual feedback as part of a compressive quality improvement program, but they recommend against use of these devices in isolation [13].

A CPR coach may assist during resuscitation to monitor and provide feedback on compression quality and coordinate tasks (e.g., switching compressors, defibrillation, intubation) [13,15,96,97]. Literature suggests that a dedicated CPR coach can reduce compression pauses while improving compression quality, compression fraction, and compression depth compliance [96,97]. A 2025 systematic review found improved compression quality, compression fraction, and guideline adherence, as well as reduced pre-shock pauses [98]. However, there was no difference in patient survival.

$EtCO_2$ can be used to monitor compressions and circulation, as $EtCO_2$ reflects pulmonary blood flow, and during cardiac arrest alveolar ventilation and metabolism remain constant [12-14,99,100]. Effective CPR increases cardiac output, leading to higher $EtCO_2$ which reflects increased perfusion. Several studies demonstrate that $EtCO_2$ values increase by 1.4 mm Hg for every 10 mm increase in compression death, with $EtCO_2$ values decreasing by 3 mm Hg for every increase in 10 breaths per minute [99-101]. $EtCO_2$ values less than 10 mm Hg during cardiac arrest suggest poor compression quality, as does a gradual decline in values [14,99,102-110]. If these are present, the compressor should be switched. $EtCO_2$ is also one of the earliest indicators of ROSC, as the return of cardiac activity results in a significant increase in $EtCO_2$. Thus, not only can $EtCO_2$ values be utilized for monitoring chest compression quality, they may also be used to predict and assess for ROSC without the need to check pulses. Systematic review data suggest values over 10 mm Hg during CPR are associated with greater likelihood of achieving ROSC, while persistent values less than 10 mm Hg after 20 min are associated with a 0.5 % likelihood of ROSC [99,104-112]. Sudden increases to normal values (30–40 mm Hg), or increases by at least 10 mm Hg, suggest ROSC [101,103,113-116]. However, there are several considerations, including medication use. This includes epinephrine, which may decrease values, and sodium bicarbonate, which may transiently increase values. However, an elevation in $EtCO_2$ with ROSC is more significant and will last longer [14,99].

Placement of an arterial line allows for continuous monitoring and can assist in resuscitation and determine ROSC, as it can directly provide the DBP and indirectly the coronary perfusion pressure. The coronary perfusion pressure is the pressure gradient of the aorta and right atrium during the decompression phase of CPR, when the myocardium receives blood flow. One study found a coronary perfusion pressure of ≥15–25 mm Hg was associated with ROSC, but this can be challenging to calculate during resuscitation [117]. Thus, utilizing DBP is a practical target. A 2016 study found physiologic monitoring with DBP was associated with higher ROSC, but there was no improvement in survival to hospital discharge or favorable neurologic outcomes [116]. Limited data suggest DBP between 25 and 40 mm Hg may be used as a target [117-120]. Thus, if a team member is present that can place an arterial line, or an arterial line is in place, DBP can be utilized to assess the effectiveness of resuscitation and monitor for ROSC. However, the team leader should not prioritize arterial line placement if there are limited personnel assisting with management.

Regional cerebral tissue oxygenation (rSO_2) utilizes noninvasive spectroscopy to detect hemoglobin [121-123]. In cardiac arrest, rSO_2 values decrease to undetectable or low levels [121]. During high-quality CPR, rSO_2 values improve, with a sudden sustained increase associated with ROSC. Mean values between 20 and 30 % are not associated with ROSC based on current literature, though there is no definitive threshold, and further study is required [122,124,125].

POCUS has demonstrated utility in several aspects of cardiac arrest management, including diagnosis of an underlying etiology, optimizing compression position, pulse checks, and prognosis [126]. Concerning CPR, POCUS may be utilized to monitor compression quality and optimize compression location. A 2020 study evaluated the feasibility of POCUS for optimizing location by assessing cardiac squeeze on POCUS and its effect on $EtCO_2$. Authors found POCUS improved compression position and $EtCO_2$ [127]. POCUS can also assess for a pulse with B-mode or color Doppler, which evaluates for pulsatility and pulsed-wave Doppler to evaluate for quantifiable pulsatile vascular flow. While a future review will discuss POCUS in greater detail, literature suggests POCUS pulse assessment of the carotid or femoral artery is rapid, reliable, and accurate when performed in the ED [128-136]. However, further data are necessary evaluating Doppler in isolation for assessment of systemic perfusion. If POCUS is utilized in cardiac arrest, the most experienced sonographer should perform the examination, and POCUS must not delay or interfere with compressions [126].

Based on the current evidence, monitoring of chest compressions is not associated with improved ROSC, survival, or neurologic outcomes, but it can provide important information and improve guideline adherence. Types of monitoring include real-time feedback, a CPR coach, $EtCO_2$ values, arterial line monitoring, rSO_2, and POCUS. If compression quality is poor, the resuscitation team must make modifications to improve CPR quality (e.g., switch compressor) in order to optimize patient outcomes.

Table 5
Monitoring CPR considerations.

- CPR quality varies significantly even when performed by trained individuals.
- There are several means of monitoring including include real-time feedback, a CPR coach, $EtCO_2$ values, arterial line monitoring, rSO_2, and POCUS.
- While literature has not demonstrated improved outcomes with CPR monitoring, any decrease in chest compression quality must be addressed to improve CPR quality.

3. Conclusions

Cardiac arrest is the loss of organized cardiac activity and systemic perfusion, which must be rapidly reversed. Treatment includes high-quality chest compressions to restore systemic perfusion. High-quality chest compressions optimize coronary and cerebral perfusion pressure, and a depth of 5–6 cm is recommended at a rate of 100–120 compressions per minute. Interruptions in compressions must be avoided, with a compression fraction of at least 60 % (over 80 % is optimal). Bystander performed compression-only CPR may be associated with improved survival to hospital discharge compared to conventional CPR with ventilations, but in trained personnel, there is likely no difference versus conventional CPR. Mechanical devices for CPR (e.g., LUCAS,

Descargado para Anonymous User (n/a) en Cayetano Heredia Pervuvian University de ClinicalKey.es por Elsevier en abril 11, 2025. Para uso personal exclusivamente. No se permiten otros usos sin autorización. Copyright ©2025. Elsevier Inc. Todos los derechos reservados.

AutoPulse) have not demonstrated an improvement in survival, ROSC, or neurologic function, but they may be used in certain situations (e.g., few rescuers available, prolonged arrest/transport). There are several means of monitoring chest compressions (real-time feedback, a CPR coach, $EtCO_2$, arterial line monitoring, rSO_2, and POCUS), which can improve guideline adherence.

CRediT authorship contribution statement

Brit Long: Writing – review & editing, Writing – original draft, Visualization, Validation, Supervision, Resources, Conceptualization. **Michael Gottlieb:** Writing – review & editing, Writing – original draft, Visualization, Validation, Supervision, Resources, Conceptualization.

Declaration of competing interest

None. No AI program was used in the construction of this manuscript.

Acknowledgements

BL and MG conceived the idea for this manuscript and contributed substantially to the writing and editing of the review. This manuscript did not utilize any grants, and it has not been presented in abstract form. This clinical review has not been published, it is not under consideration for publication elsewhere, its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. This review does not reflect the views or opinions of the U.S. government, Department of Defense, Defense Health Agency, Brooke Army Medical Center, the U.S. Army, U.S. Air Force, or SAUSHEC EM Residency Program.

References

[1] Sayre MR, Koster RW, Botha M, et al. Part 5: adult basic life support: 2010 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. Circulation. 2010;122(16 Suppl 2):S298–324.
[2] Berg KM, Bray JE, Ng KC, et al, Collaborators. 2023 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations: summary from the basic life support; advanced life support; pediatric life support; neonatal life support; education, implementation, and teams; and first aid task forces. Circulation. 2023 Dec 12;148(24):e187–280. https://doi.org/10.1161/CIR.0000000000001179. Epub 2023 Nov 9. Erratum in: Circulation. 2024 Apr 16;149(16):e1128. doi: 10.1161/CIR.0000000000001242. Erratum in: Circulation. 2024 Jun 11;149(24):e1411.
[3] Tsao CW, Aday AW, Almarzooq ZI, et al. Heart disease and stroke statistics-2022 update: a report from the American Heart Association. Circulation. 2022 Feb 22;145(8):e153–639.
[4] Writing Group Members, Roger VL, Go AS, et al. Heart disease and stroke statistics – 2012 update: a report from the American Heart Association. Circulation. 2012 (125):e2-20.
[5] WRITING GROUP MEMBERS, Lloyd-Jones D, Adams RJ, American Heart Association Statistics Committee and Stroke Statistics Subcommittee, et al. Heart disease and stroke statistics–2010 update: a report from the American Heart Association. Circulation. 2010 Feb 23;121(7):e46-215.
[6] Kong MH, Fonarow GC, Peterson ED, et al. Systematic review of the incidence of sudden cardiac death in the United States. J Am Coll Cardiol. 2011 Feb 15;57(7):794–801.
[7] Herlitz J, Svensson L, Engdahl J, et al. Characteristics of cardiac arrest and resuscitation by age group: an analysis from the Swedish cardiac arrest registry. Am J Emerg Med. 2007;25:1025–31.
[8] Ong ME, Stiell I, Osmond MH, et al. Etiology of pediatric out-of-hospital cardiac arrest by coroner's diagnosis. Resuscitation. 2006;68:335–42.
[9] Wyckoff MH, Singletary EM, Soar J, COVID-19 Working Group, et al. 2021 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations: summary from the basic life support; advanced life support; neonatal life support; education, implementation, and teams; first aid task forces; and the COVID-19 working group. Resuscitation. 2021 Dec;169:229–311.
[10] Wyckoff MH, Singletary EM, Soar J, et al, Collaborators. 2021 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations: summary from the basic life support; advanced life support; neonatal life support; education, implementation, and teams; first aid task forces; and the COVID-19 working group. Circulation. 2022 Mar;145(9):e645–721.
[11] Panchal AR, Bartos JA, Cabañas JG, et al. Adult basic and advanced life support writing group. Part 3: adult basic and advanced life support: 2020 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. Circulation. 2020 Oct 20;142(16_suppl_2):S366–468.
[12] Berg RA, Hemphill R, Abella BS, et al. Part 5: adult basic life support: 2010 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. Circulation. 2010 Nov 2;122(18 Suppl 3):S685–705.
[13] Greif R, Bray JE, Djärv T, et al. 2024 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations: summary from the basic life support; advanced life support; pediatric life support; neonatal life support; education, implementation, and teams; and first aid task forces. Circulation. 2024 Dec 10;150(24):e580–687.
[14] Long B, Gottlieb M. Cardiopulmonary resuscitation: the importance of the basics. Emerg Med Clin North Am. 2023 Aug;41(3):509–28.
[15] Olasveengen TM, Mancini ME, Perkins GD, et al, Adult Basic Life Support Collaborators. Adult basic life support: 2020 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. Circulation. 2020 Oct 20;142(16_suppl_1):S41–91.
[16] Ochoa FJ, Ramalle-Gómara E, Carpintero JM, et al. Competence of health professionals to check the carotid pulse. Resuscitation. 1998;37:173–5.
[17] Perkins GD, Stephenson B, Hulme J, Monsieurs KG. Birmingham assessment of breathing study (BABS). Resuscitation. 2005;64:109–13.
[18] Van Hoeyweghen RJ, Bossaert LL, Mullie A, et al. Quality and efficiency of bystander CPR. Belg Cereb Resuscitation Study Group Resuscitation. 1993 Aug;26(1):47–52.
[19] Gallagher EJ, Lombardi G, Gennis P. Effectiveness of bystander cardiopulmonary resuscitation and survival following out-of-hospital cardiac arrest. JAMA. 1995 Dec 27;274(24):1922–5.
[20] Berg RA, Sanders AB, Kern KB, et al. Adverse hemodynamic effects of interrupting chest compressions for rescue breathing during cardiopulmonary resuscitation for ventricular fibrillation cardiac arrest. Circulation. 2001 Nov 13;104(20):2465–70.
[21] Kellum MJ, Kennedy KW, Barney R, et al. Cardiocerebral resuscitation improves neurologically intact survival of patients with out-of-hospital cardiac arrest. Ann Emerg Med. 2008 Sep;52(3):244–52.
[22] Abella BS, Sandbo N, Vassilatos P, et al. Chest compression rates during cardiopulmonary resuscitation are suboptimal: a prospective study during in-hospital cardiac arrest. Circulation. 2005 Feb 1;111(4):428–34.
[23] Perkins GD, Kocierz L, Smith SC, et al. Compression feedback devices over estimate chest compression depth when performed on a bed. Resuscitation. 2009 Jan;80(1):79–82.
[24] Feneley MP, Maier GW, Kern KB, et al. Influence of compression rate on initial success of resuscitation and 24 hour survival after prolonged manual cardiopulmonary resuscitation in dogs. Circulation. 1988 Jan;77(1):240–50.
[25] Stiell IG, Brown SP, Christenson J, Resuscitation Outcomes Consortium (ROC) Investigators, et al. What is the role of chest compression depth during out-of-hospital cardiac arrest resuscitation? Crit Care Med. 2012;40:1192–8.
[26] Stiell IG, Brown SP, Nichol G, Resuscitation Outcomes Consortium Investigators, et al. What is the optimal chest compression depth during out-of-hospital cardiac arrest resuscitation of adult patients? Circulation. 2014;130:1962–70.
[27] SOS-KANTO study group. Cardiopulmonary resuscitation by bystanders with chest compression only (SOS-KANTO): an observational study. Lancet. 2007 Mar 17;369(9565):920–6.
[28] Edelson DP, Abella BS, Kramer-Johansen J, et al. Effects of compression depth and pre-shock pauses predict defibrillation failure during cardiac arrest. Resuscitation. 2006;71:137–45.
[29] Babbs CF, Kemeny AE, Quan W, Freeman G. A new paradigm for human resuscitation research using intelligent devices. Resuscitation. 2008;77:306–15.
[30] Yannopoulos D, McKnite S, Aufderheide TP, et al. Effects of incomplete chest wall decompression during cardiopulmonary resuscitation on coronary and cerebral perfusion pressures in a porcine model of cardiac arrest. Resuscitation. 2005 Mar;64(3):363–72.
[31] Nishisaki A, Maltese MR, Niles DE, et al. Backboards are important when chest compressions are provided on a soft mattress. Resuscitation. 2012;83:1013–20.
[32] Noordergraaf GJ, Paulussen IW, Venema A, et al. The impact of compliant surfaces on in-hospital chest compressions: effects of common mattresses and a backboard. Resuscitation. 2009;80:546–52.
[33] Dewan M, Schachna E, Eastwood K, Perkins G, Bray J, International Liaison Committee on Resuscitation Basic Life Support Task Force. The optimal surface for delivery of CPR: an updated systematic review and meta-analysis. Resusc Plus. 2024 Jul 20(19):100718.
[34] Shin J, Rhee JE, Kim K. Is the inter-nipple line the correct hand position for effective chest compression in adult cardiopulmonary resuscitation? Resuscitation. 2007;75:305–10.
[35] Anez C, Becerra-Bolaños Á, Vives-Lopez A, Rodríguez-Pérez A. Cardiopulmonary resuscitation in the prone position in the operating room or in the intensive care unit: a systematic review. Anesth Analg. 2021 Feb 1;132(2):285–92.
[36] Christenson J, Andrusiek D, Everson-Stewart S, Resuscitation Outcomes Consortium Investigators, et al. Chest compression fraction determines survival in patients with out-of-hospital ventricular fibrillation. Circulation. 2009 Sep 29;120(13):1241–7.
[37] Kern KB, Hilwig RW, Berg RA, et al. Importance of continuous chest compressions during cardiopulmonary resuscitation: improved outcome during a simulated single lay-rescuer scenario. Circulation. 2002 Feb 5;105(5):645–9.

Descargado para Anonymous User (n/a) en Cayetano Heredia Pervuvian University de ClinicalKey.es por Elsevier en abril 11, 2025. Para uso personal exclusivamente. No se permiten otros usos sin autorización. Copyright ©2025. Elsevier Inc. Todos los derechos reservados.

[38] Eftestøl T, Sunde K, Steen PA. Effects of interrupting precordial compressions on the calculated probability of defibrillation success during out-of-hospital cardiac arrest. Circulation. 2002 May 14;105(19):2270–3.
[39] Garza AG, Gratton MC, Salomone JA, Lindholm D, McElroy J, Archer R. Improved patient survival using a modified resuscitation protocol for out-of-hospital cardiac arrest. Circulation. 2009 May 19;119(19):2597–605.
[40] Sayre MR, Cantrell SA, White LJ, et al. Impact of the 2005 American Heart Association cardiopulmonary resuscitation and emergency cardiovascular care guidelines on out-of-hospital cardiac arrest survival. Prehosp Emerg Care. 2009.;Oct-Dec;13(4):469-77.
[41] Cheskes S, Schmicker RH, Christenson J, et al. Resuscitation outcomes consortium (ROC) investigators. Perishock pause: an independent predictor of survival from out-of-hospital shockable cardiac arrest. Circulation. 2011 Jul 5;124(1):58–66.
[42] Brouwer TF, Walker RG, Chapman FW, Koster RW. Association between chest compression interruptions and clinical outcomes of ventricular fibrillation out-of-hospital cardiac arrest. Circulation. 2015 Sep 15;132(11):1030–7.
[43] Dewolf P, Wauters L, Clarebout G, et al. Assessment of chest compression interruptions during advanced cardiac life support. Resuscitation. 2021 Aug;165:140–7.
[44] Andreka P, Frenneaux MP. Haemodynamics of cardiac arrest and resuscitation. Curr Opin Crit Care. 2006 Jun;12(3):198–203.
[45] Ewy GA. Cardiocerebral resuscitation: the new cardiopulmonary resuscitation. Circulation. 2005 Apr 26;111(16):2134–42.
[46] Nichol G, Leroux B, Wang H, ROC Investigators, et al. Trial of continuous or interrupted chest compressions during CPR. N Engl J Med. 2015 Dec 3;373(23):2203–14.
[47] Rea TD, Helbock M, Perry S, et al. Increasing use of cardiopulmonary resuscitation during out-of-hospital ventricular fibrillation arrest: survival implications of guideline changes. Circulation. 2006;114:2760–5.
[48] Lloyd MS, Heeke B, Walter PF. Hands-on defibrillation: an analysis of electrical current flow through rescuers in direct contact with patients during biphasic external defibrillation. Circulation. 2008;117(19):2510–4.
[49] Iwami T, Kitamura T, Kiyohara K, Kawamura T. Dissemination of chest compression-only cardiopulmonary resuscitation and survival after out-of-hospital cardiac arrest. Circulation. 2015;132:415–22.
[50] Kitamura T, Iwami T, Kawamura T, et al. Implementation working Group for all-Japan Utstein Registry of the fire and disaster management agency. Bystander-initiated rescue breathing for out-of-hospital cardiac arrests of noncardiac origin. Circulation. 2010;122:293–9.
[51] Ogawa T, Akahane M, Koike S, et al. Outcomes of chest compression only CPR versus conventional CPR conducted by lay people in patients with out of hospital cardiopulmonary arrest witnessed by bystanders: nationwide population based observational study. BMJ. 2011;342:c7106.
[52] Svensson L, Bohm K, Castrèn M, et al. Compression-only CPR or standard CPR in out-of-hospital cardiac arrest. N Engl J Med. 2010;363:434–42.
[53] Rea TD, Fahrenbruch C, Culley L, et al. CPR with chest compression alone or with rescue breathing. N Engl J Med. 2010;363:423–33.
[54] Kern KB, Ewy GA, Voorhees WD, et al. Myocardial perfusion pressure: a predictor of 24-hour survival during prolonged cardiac arrest in dogs. Resuscitation. 1998;16:241–50.
[55] Ashoor HM, Lillie E, Zarin W, ILCOR Basic Life Support Task Force, et al. Effectiveness of different compression-to-ventilation methods for cardiopulmonary resuscitation: a systematic review. Resuscitation. 2017 Sep;118:112–25.
[56] Mader TJ, Paquette AT, Salcido DD, et al. The effect of the preshock pause on coronary perfusion pressure decay and rescue shock outcome in porcine ventricular fibrillation. Prehosp Emerg Care. 2009;13:487–94.
[57] Zhan L, Yang LJ, Huang Y, He Q, Liu GJ. Continuous chest compression versus interrupted chest compression for cardiopulmonary resuscitation of non-asphyxial out-of-hospital cardiac arrest. Cochrane Database Syst Rev. 2017 Mar 27;3(3):CD010134.
[58] Hüpfl M, Selig HF, Nagele P. Chest-compression-only versus standard cardiopulmonary resuscitation: a meta-analysis. Lancet. 2010 Nov 6;376(9752):1552–7.
[59] Olasveengen TM, de Caen AR, Mancini ME, et al, ILCOR Collaborators. 2017 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations summary. Resuscitation. 2017 Dec;121:201–14.
[60] Rea TD, Fahrenbruch C, Culley L, et al. CPR with chest compression alone or with rescue breathing. N Engl J Med. 2010 Jul 29;363(5):423–33.
[61] Hallstrom A, Cobb L, Johnson E, Copass M. Cardiopulmonary resuscitation by chest compression alone or with mouth-to-mouth ventilation. N Engl J Med. 2000 May 25;342(21):1546–53.
[62] Yao L, Wang P, Zhou L, et al. Compression-only cardiopulmonary resuscitation vs standard cardiopulmonary resuscitation: an updated meta-analysis of observational studies. Am J Emerg Med. 2014;32(6):517–23.
[63] Ivan I, Budiman F, Ruby R, et al. Current evidence of survival benefit between chest-compression only versus standard cardiopulmonary resuscitation in out-of-hospital cardiac arrest : updated systematic review and meta-analysis of randomized controlled trials with trial sequential analysis. Herz. 2021 Sep;46(Suppl. 2):198–208.
[64] Bielski K, Smereka J, Chmielewski J, et al. Meta-analysis of chest compression-only versus conventional cardiopulmonary resuscitation by bystanders for adult with out-of-hospital cardiac arrest. Cardiol J. 2023;30(4):606–13.
[65] Sun M, Zhu A, Tang Y. Continuous compression with asynchronous ventilation improves CPR prognosis? A meta-analysis from human and animal studies. Am J Emerg Med. 2023 Feb;64:26–36.
[66] AutoPulse resuscitation system. Zoll. Available at. https://www.zoll.com/products/automated-cpr/autopulse-for-ems. [Accessed 21 August 2022].
[67] LUCAS chest compression system. Stryker. Available at. https://www.lucas-cpr.com. [Accessed 21 August 2022].
[68] Poole K, Couper K, Smyth MA, et al. Mechanical CPR: who? When? How? Crit Care. 2018 May 29;22(1):140.
[69] Sheraton M, Columbus J, Surani S, et al. Effectiveness of mechanical chest compression devices over manual cardiopulmonary resuscitation: a systematic review with meta-analysis and trial sequential analysis. West J Emerg Med. 2021 Jul 19;22(4):810–9.
[70] Wang PL, Brooks SC. Mechanical versus manual chest compressions for cardiac arrest. Cochrane Database Syst Rev. 2018 Aug 20;8(8):CD007260.
[71] Zhu N, Chen Q, Jiang Z, et al. A meta-analysis of the resuscitative effects of mechanical and manual chest compression in out-of-hospital cardiac arrest patients. Crit Care. 2019;23:100.
[72] Liu M, Shuai Z, Ai J, et al. Mechanical chest compression with LUCAS device does not improve clinical outcome in out-of-hospital cardiac arrest patients: a systematic review and meta-analysis. Med (Baltim). 2019;98:e17550.
[73] Gao Y, Sun T, Yuan D, et al. Safety of mechanical and manual chest compressions in cardiac arrest patients: a systematic review and meta-analysis. Resuscitation. 2021;169:124–35.
[74] Larik MO, Ahmed A, Shiraz MI, Shiraz SA, Anjum MU, Bhattarai P. Comparison of manual chest compression versus mechanical chest compression for out-of-hospital cardiac arrest: a systematic review and meta-analysis. Med (Baltim). 2024 Feb 23;103(8):e37294.
[75] Esibov A, Banville I, Chapman FW, et al. Mechanical chest compressions improved aspects of CPR in the LINC trial. Resuscitation. 2015;91:116–21.
[76] Yost D, Phillips RH, Gonzales L, et al. Assessment of CPR interruptions from transthoracic impedance during use of the LUCASÔ mechanical chest compression system. Resuscitation. 2012;83(8):961–5.
[77] Levy M, Yost D, Walker RG, et al. A quality improvement initiative to optimize use of a mechanical chest compression device within a high-performance CPR approach to out-of-hospital cardiac arrest resuscitation. Resuscitation. 2015;92:32–7.
[78] Lindner KH, Pfenninger EG, Lurie KG, et al. Effects of active compression-decompression resuscitation on myocardial and cerebral blood flow in pigs. Circulation. 1993 Sep;88(3):1254–63.
[79] Tucker KJ, Redberg RF, Schiller NB, Cohen TJ. Active compression-decompression resuscitation: analysis of transmitral flow and left ventricular volume by transesophageal echocardiography in humans. Cardiopulmonary Resuscitation Working Group. J Am Coll Cardiol. 1993 Nov 1;22(5):1485–93.
[80] Cohen TJ, Goldner BG, Maccaro PC, et al. A comparison of active compression-decompression cardiopulmonary resuscitation with standard cardiopulmonary resuscitation for cardiac arrests occurring in the hospital. N Engl J Med. 1993 Dec 23;329(26):1918–21.
[81] Lafuente-Lafuente C, Melero-Bascones M. Active chest compression-decompression for cardiopulmonary resuscitation. Cochrane Database Syst Rev. 2013 Sep 20;2013(9):CD002751.
[82] Plaisance P, Soleil C, Lurie KG, et al. Use of an inspiratory impedance threshold device on a facemask and endotracheal tube to reduce intrathoracic pressures during the decompression phase of active compression-decompression cardiopulmonary resuscitation. Crit Care Med. 2005 May;33(5):990–4.
[83] Plaisance P, Lurie KG, Vicaut E, et al. Evaluation of an impedance threshold device in patients receiving active compression-decompression cardiopulmonary resuscitation for out of hospital cardiac arrest. Resuscitation. 2004 Jun;61(3):265–71.
[84] Aufderheide TP, Nichol G, Rea TD, Resuscitation Outcomes Consortium (ROC) Investigators, et al. A trial of an impedance threshold device in out-of-hospital cardiac arrest. N Engl J Med. 2011 Sep 1;365(9):798–806.
[85] Aufderheide TP, Pirrallo RG, Provo TA, et al. Clinical evaluation of an inspiratory impedance threshold device during standard cardiopulmonary resuscitation in patients with out-of-hospital cardiac arrest. Crit Care Med. 2005;33(4):734–40.
[86] Lurie KG, Coffeen P, Shultz J, et al. Improving active compression-decompression cardiopulmonary resuscitation with an inspiratory impedance valve. Circulation. 1995;91(6):1629–32.
[87] Lurie KG, Lindner KH. Recent advances in cardiopulmonary resuscitation. J Cardiovasc Electrophysiol. 1997;8(5):584–600.
[88] Yannopoulos D, Aufderheide TP, Abella BS, et al. Quality of CPR: an important effect modifier in cardiac arrest clinical outcomes and intervention effectiveness trials. Resuscitation. 2015;94:106–13.
[89] Aufderheide TP, Frascone RJ, Wayne MA, et al. Treatment of out-of-hospital cardiac arrest with an impedance threshold device and active compression decompression cardiopulmonary resuscitation improves survival with good neurological outcome: results from the resqtrial. Resuscitation. 2010;81(2):S21.
[90] Pepe PE, Scheppke KA, Antevy PM, et al. Confirming the clinical safety and feasibility of a bundled methodology to improve cardiopulmonary resuscitation involving a head-up/torso-up chest compression technique. Crit Care Med. 2019 Mar;47(3):449–55.
[91] Moore JC, Duval S, Lick C, et al. Faster time to automated elevation of the head and thorax during cardiopulmonary resuscitation increases the probability of return of spontaneous circulation. Resuscitation. 2022 Jan;170:63–9.
[92] Moore JC, Pepe PE, Scheppke KA, et al. Head and thorax elevation during cardiopulmonary resuscitation using circulatory adjuncts is associated with improved survival. Resuscitation. 2022 Oct;179:9–17.
[93] Duval S, Pepe PE, Aufderheide TP, et al. Optimal combination of compression rate and depth during cardiopulmonary resuscitation for functionally favorable survival. JAMA Cardiol. 2019 Sep 1;4(9):900–8.
[94] Lyngby RM, Quinn T, Oelrich RM, et al. Association of real-time feedback and cardiopulmonary-resuscitation quality delivered by ambulance personnel for out-of-hospital cardiac arrest. J Am Heart Assoc. 2023 Oct 17;12(20):e029457.

Descargado para Anonymous User (n/a) en Cayetano Heredia Pervuvian University de ClinicalKey.es por Elsevier en abril 11, 2025. Para uso personal exclusivamente. No se permiten otros usos sin autorización. Copyright ©2025. Elsevier Inc. Todos los derechos reservados.

[95] Masterson S, Norii T, Yabuki M, BLS ILCOR Task Force, et al. Real-time feedback for CPR quality - a scoping review. Resusc Plus. 2024 Jul 31(19):100730.
[96] Goharani R, Vahedian-Azimi A, Farzanegan B, et al, MORZAK Collaborative. Real-time compression feedback for patients with in-hospital cardiac arrest: a multicenter randomized controlled clinical trial. J Intensive Care. 2019 Jan 22(7):5.
[97] Cheng A, Duff JP, Kessler D, International Network for Simulation-based Pediatric Innovation Research and Education (INSPIRE) CPR, et al. Optimizing CPR performance with CPR coaching for pediatric cardiac arrest: a randomized simulation-based clinical trial. Resuscitation. 2018 Nov;132:33–40.
[98] Lauridsen KG, Bürgstein E, Nabecker S, et al. Cardiopulmonary resuscitation coaching for resuscitation teams: a systematic review. Resusc Plus. 2025 Jan 8 (21):100868.
[99] Long B, Koyfman A, Vivirito MA. Capnography in the emergency department: a review of uses, waveforms, and limitations. J Emerg Med. 2017 Dec;53(6):829–42.
[100] Nassar BS, Schmidt GA. Capnography during critical illness. Chest. 2016 Feb;149 (2):576–85.
[101] Sheak KR, Wiebe DJ, Leary M, et al. Quantitative relationship between end-tidal carbon dioxide and CPR quality during both in-hospital and out-of-hospital cardiac arrest. Resuscitation. 2015;89:149–54.
[102] Paiva EF, Paxton JH, O'Neil BJ. The use of end-tidal carbon dioxide (ETCO2) measurement to guide management of cardiac arrest: a systematic review. Resuscitation. 2018;123:1–7.
[103] Neumar RW, Shuster M, Callaway CW, et al. Part 1: executive summary: 2015 American Heart Association guidelines update for cardiopulmonary resuscitation and emergency cardiovascular care. Circulation. 2015;132(18 Suppl 2):S315–67.
[104] Davis DP, Sell RE, Wilkes N, et al. Electrical and mechanical recovery of cardiac function following out-of-hospital cardiac arrest. Resuscitation. 2013;84:25.
[105] Sanders AB, Kern KB, Otto CW, et al. End-tidal carbon dioxide monitoring during cardiopulmonary resuscitation. A prognostic indicator for survival. JAMA. 1989; 262:1347.
[106] Chen J-J, Lee Y-K, Hou S-W, et al. End-tidal carbon dioxide monitoring may be associated with a higher possibility of return of spontaneous circulation during out-of-hospital cardiac arrest: a population-based study. Scand J Trauma Resusc Emerg Med. 2015;23:104.
[107] Murphy RA, Bobrow BJ, Spaite DW, et al. Association between prehospital CPR quality and end-tidal carbon dioxide levels in out-of-hospital cardiac arrest. Prehosp Emerg Care. 2016 May-Jun.;20(3):369-77.
[108] Levine RL, Wayne MA, Miller CC. End-tidal carbon dioxide and outcome of out-of-hospital cardiac arrest. N Engl J Med. 1997;337:301.
[109] Wayne MA, Levine RL, Miller CC. Use of end-tidal carbon dioxide to predict outcome in prehospital cardiac arrest. Ann Emerg Med. 1995;25:762.
[110] Grmec S, Klemen P. Does the end-tidal carbon dioxide (EtCO2) concentration have prognostic value during out-of-hospital cardiac arrest? Eur J Emerg Med. 2001;8: 263.
[111] Steedman DJ, Robertson CE. Measurement of end-tidal carbon dioxide concentration during cardiopulmonary resuscitation. Arch Emerg Med. 1990;7(3):129–34.
[112] Einav S, Bromiker R, Weiniger CF, Matot I. Mathematical modeling for prediction of survival from resuscitation based on computerized continuous capnography: proof of concept. Acad Emerg Med. 2011;18:468–75.
[113] Eckstein M, Hatch L, Malleck J, et al. End-tidal CO2 as a predictor of survival in out-of-hospital cardiac arrest. Resuscitation. 2016 Jul;104:53–8.
[114] Falk JL, Rackow EC, Weil MH. End-tidal carbon dioxide concentration during cardiopulmonary resuscitation. N Engl J Med. 1988;318:607.
[115] Garnett AR, Ornato JP, Gonzalez ER, Johnson EB. End-tidal carbon dioxide monitoring during cardiopulmonary resuscitation. JAMA. 1987;257:512.
[116] Sutton RM, French B, Meaney PA, et al. Physiologic monitoring of CPR quality during adult cardiac arrest: a propensity-matched cohort study. Resuscitation. 2016; 106:76–82.
[117] Paradis NA, Martin GB, Rivers EP, et al. Coronary perfusion pressure and the return of spontaneous circulation in human cardiopulmonary resuscitation. JAMA. 1990; 263(8):1106–13.
[118] Sutton RM, Friess SH, Maltese MR, et al. Hemodynamic-directed cardiopulmonary resuscitation during in-hospital cardiac arrest. Resuscitation. 2014;85(8):983–6.
[119] Drumheller BC, Pinizzotto J, Overberger RC, et al. Goal-directed cardiopulmonary resuscitation for refractory out-of-hospital cardiac arrest in the emergency department: a feasibility study. Resusc Plus. 2021;7:100159.
[120] Meaney PA, Bobrow BJ, Mancini ME, et al. Cardiopulmonary resuscitation quality: [corrected] improving cardiac resuscitation outcomes both inside and outside the hospital: a consensus statement from the American Heart Association. Circulation. 2013;128(4):417–35.
[121] Prosen G, Strnad M, Doniger SJ, et al. Cerebral tissue oximetry levels during prehospital management of cardiac arrest - a prospective observational study. Resuscitation. 2018;129:141–5.
[122] Schnaubelt S, Sulzgruber P, Menger J, et al. Regional cerebral oxygen saturation during cardiopulmonary resuscitation as a predictor of return of spontaneous circulation and favourable neurological outcome - a review of the current literature. Resuscitation. 2018;125:39–47.
[123] Takegawa R, Shiozaki T, Ogawa Y, et al. Usefulness of cerebral rSO. Resuscitation. 2019;139:201–7.
[124] Cournoyer A, Iseppon M, Chauny JM, et al. Near-infrared spectroscopy monitoring during cardiac arrest: a systematic review and meta-analysis. Acad Emerg Med. 2016;23(8):851–62.
[125] Liu Y, Jing K, Liu H, et al. Association between cerebral oximetry and return of spontaneous circulation following cardiac arrest: a systematic review and meta-analysis. PloS One. 2020;15(8):e0234979.
[126] Gottlieb M, Alerhand S. Managing cardiac arrest using ultrasound. Ann Emerg Med. 2023 May;81(5):532–42.
[127] Zanatta M, Lorenzi C, Scorpiniti M, et al. Ultrasound-guided chest compressions in out-of-hospital cardiac arrests. J Emerg Med. 2020 Dec;59(6):e225–33.
[128] Adedipe AA, Fly DL, Schwitz SD, et al. Carotid Doppler blood flow measurement during cardiopulmonary resuscitation is feasible: a first in man study. Resuscitation. 2015;96:121–5.
[129] Germanoska B, Coady M, Ng S, et al. The reliability of carotid ultrasound in determining the return of pulsatile ow: a pilot study. Ultrasound. 2018;26:118–26.
[130] Badra K, Coutin A, Simard R, et al. The POCUS pulse check: a randomized controlled crossover study comparing pulse detection by palpation versus by point-of-care ultrasound. Resuscitation. 2019;139:17–23.
[131] Simard RD, Unger AG, Betz M, et al. The POCUS pulse check: a case series on a novel method for determining the presence of a pulse using point-of-care ultrasound. J Emerg Med. 2019;56:674–9.
[132] Sanchez S, Miller M, Asha S. Assessing the validity of two- dimensional carotid ultrasound to detect the presence and absence of a pulse. Resuscitation. 2020;157: 67–73.
[133] Smith DJ, Simard R, Chenkin J. Checking the pulse in the 21st century: interobserver reliability of carotid pulse detection by point-of-care ultrasound. Am J Emerg Med. 2021;45:280–3.
[134] Schwartz BE, Gandhi P, Najafali D, et al. Manual palpation vs. femoral arterial Doppler ultrasound for comparison of pulse check time during cardiopulmonary resuscitation in the emergency department: a pilot study. J Emerg Med. 2021;61: 720–30.
[135] Kang SY, Jo IJ, Lee G, et al. Point-of-care ultrasound compression of the carotid artery for pulse determination in cardiopulmonary resuscitation. Resuscitation. 2022;S0300-9572:00590–1.
[136] Cohen AL, Li T, Becker LB, et al. Femoral artery Doppler ultrasound is more accurate than manual palpation for pulse detection in cardiac arrest. Resuscitation. 2022; 173:156–65.

Descargado para Anonymous User (n/a) en Cayetano Heredia Pervuvian University de ClinicalKey.es por Elsevier en abril 11, 2025. Para uso personal exclusivamente. No se permiten otros usos sin autorización. Copyright ©2025. Elsevier Inc. Todos los derechos reservados.